



DOCKET NO.: 211710US0X

TITLE OF THE INVENTION

Nucleotide sequences which code for the metE gene

BACKGROUND OF THE INVENTION

5 Field of the Invention

The invention provides nucleotide sequences from coryneform bacteria which code for the metE gene and a process for the fermentative preparation of amino acids, in particular L-methionine, using bacteria in which the metE gene is enhanced.

10 Description of the Related Art

L-Amino acids, in particular L-methionine, are used in human medicine and in the pharmaceuticals industry, in the foodstuffs industry and very particularly in animal nutrition.

15 It is known that amino acids are prepared by fermentation from strains of coryneform bacteria, in particular Corynebacterium glutamicum. Because of their great importance, work is constantly being undertaken to improve the preparation process. Improvements to the process can relate to fermentation measures stirring and supply of oxygen, or to the composition of the nutrient media such as the sugar
20 concentration during the fermentation, or to the working up of the product by, for example, ion exchange chromatography, or to the intrinsic output properties of the microorganism itself.

25 Methods of mutagenesis, selection and mutant selection are used to improve the output properties of these microorganisms. Strains which are resistant to antimetabolites, such as e.g. the methionine analogue α -methyl-methionine, ethionine, norleucine, N-acetylnorleucine, S-trifluoromethylhomocysteine,
30 2-amino-5-heprenoitic acid, seleno-methionine, methionine-sulfoximine, methoxine, 1-aminocyclopentane-carboxylic acid, or are auxotrophic for metabolites of regulatory importance



and produce amino acids, such as e.g. L-methionine, are obtained in this manner.

Recombinant DNA techniques have also been employed for some years for improving the Corynebacterium strains which produce L-amino acids, by amplifying individual amino acid biosynthesis genes and investigating their effect on the amino acid production.

SUMMARY OF THE INVENTION

An object of the present invention is to provide new measures for improved fermentative preparation of amino acids, in particular L-methionine.

When L-methionine or methionine are mentioned in the following, the salts, such as methionine hydrochloride or methionine sulfate are also meant.

The invention provides an isolated polynucleotide from coryneform bacteria, comprising a polynucleotide sequence which codes for the metE gene, chosen from the group consisting of

- a) polynucleotide which is at least 70% identical to a polynucleotide that codes for a polypeptide which comprises the amino acid sequence of SEQ ID No. 2,
 - b) polynucleotide which codes for a polypeptide that comprises an amino acid sequence which is at least 70% identical to the amino acid sequence of SEQ ID No. 2,
 - c) polynucleotide which is complementary to the polynucleotides of a) or b), and
 - d) polynucleotide comprising at least 15 successive nucleotides of the polynucleotide sequence of a), b) or c),
- and the corresponding polypeptide having the enzymatic activity of homocysteine methyltransferase I.

The invention also provides the above-mentioned polynucleotides as DNA which is capable of replication, comprising:

- (i) the nucleotide sequence shown in SEQ ID No. 1, or
- 5 (ii) at least one sequence which corresponds to sequence (i) within the range of the degeneration of the genetic code, or
- (iii) at least one sequence which hybridizes with the sequence complementary to sequence (i) or (ii), and
- 10 (iv) sense mutations of neutral function in (i).

The invention also provides

- a polynucleotide comprising the nucleotide sequence as shown in SEQ ID No. 1;
- 15 a polynucleotide that codes for a polypeptide which comprises the amino acid sequence as shown in SEQ ID No. 2;
- a vector containing the polynucleotide according to the invention, in particular a shuttle vector or plasmid vector, and
- 20 and coryneform bacteria serving as the host cell, which contain the vector or in which the metE gene is enhanced.

The invention also provides polynucleotides which are obtained by screening a corresponding gene library, which comprises the complete gene having the polynucleotide sequence corresponding to SEQ ID No. 1, by means of hybridization with a probe which

25 comprises the sequence of the polynucleotide mentioned, according to SEQ ID No. 1 or a fragment thereof, and isolation of the DNA sequence mentioned.

BRIEF DESCRIPTION OF THE FIGURES

30 Fig. 1 shows plasmid pCREmetAE.

Fig. 2 shows plasmid pCREmetAEY.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

Polynucleotides which comprise the sequences according to the invention are suitable as hybridization probes for RNA, cDNA and DNA, in order to isolate, in the full length, nucleic acids or polynucleotides or genes which code for homocysteine methyltransferase I or to isolate those nucleic acids or polynucleotides or genes which have a high similarity of sequence or homology with that of the homocysteine methyltransferase I gene.

Polynucleotides according to the invention are furthermore suitable as primers with which the DNA of genes that code for homocysteine methyltransferase I can be prepared by the polymerase chain reaction (PCR).

Such oligonucleotides that serve as probes or primers comprise at least 30, preferably at least 20, very particularly at least 15 successive nucleotides. Oligonucleotides which have a length of at least 40 or 50 nucleotides are also suitable. Oligonucleotides with a length of at least 100, 150, 200, 250 or 300 nucleotides are optionally also suitable.

"Isolated" means separated out of its natural environment.

"Polynucleotide" in general relates to polyribonucleotides and polydeoxyribonucleotides, it being possible for these to be non-modified RNA or DNA or modified RNA or DNA.

"Polypeptides" are understood as meaning peptides or proteins which comprise two or more amino acids bonded via peptide bonds.

The polypeptides according to the invention include a polypeptide according to SEQ ID No. 2, in particular those with the biological activity of homocysteine methyltransferase I, and also those which are at least 70%, preferably at least 80% and in particular at least 90% to 95% identical to the

polypeptide according to SEQ ID No. 2 and have the activity mentioned.

The invention moreover provides a process for the fermentative preparation of amino acids, in particular L-methionine, using coryneform bacteria which in particular already produce amino acids, and in which the nucleotide sequences which code for the metE gene are enhanced, in particular over-expressed.

The term "enhancement" in this connection describes an increase in the intracellular activity of one or more enzymes (proteins) in a microorganism which are coded by the corresponding DNA, for example by increasing the number of copies of the gene or genes, using a potent promoter or using a gene or allele which codes for a corresponding enzyme (protein) having a high activity, and optionally combining these measures.

By enhancement measures, in particular over-expression, the activity or concentration of the corresponding protein is in general increased by at least 10%, 25%, 50%, 75%, 100%, 150%, 200%, 300%, 400% or 500%, up to a maximum of 1000% or 2000%, based on the starting microorganism.

The microorganisms which the present invention provides can prepare L-amino acids, in particular L-methionine, from glucose, sucrose, lactose, fructose, maltose, molasses, starch, cellulose or from glycerol and ethanol. They can be representatives of coryneform bacteria, in particular of the genus *Corynebacterium*. Of the genus *Corynebacterium*, there may be mentioned in particular the species *Corynebacterium glutamicum*, which is known among experts for its ability to produce L-amino acids.

Suitable strains of the genus *Corynebacterium*, in particular of the species *Corynebacterium glutamicum* (*C. glutamicum*), are in particular the known wild-type strains

Corynebacterium glutamicum ATCC13032

Corynebacterium acetoglutamicum ATCC15806

Corynebacterium acetoacidophilum ATCC13870
Corynebacterium thermoaminogenes FERM BP-1539
Corynebacterium melassecola ATCC17965
Brevibacterium flavum ATCC14067
5 Brevibacterium lactofermentum ATCC13869 and
Brevibacterium divaricatum ATCC14020

or L-amino acid-producing mutants or strains prepared
therefrom, such as, for example, the L-methionine-producing
strain

10 Corynebacterium glutamicum ATCC21608.

The new metE gene from C. glutamicum which codes for the
enzyme homocysteine methyltransferase I (EC 2.1.1.14) has been
isolated.

To isolate the metE gene or also other genes of C. glutamicum,
a gene library of this microorganism is first set up in
Escherichia coli (E. coli). The setting up of gene libraries
is described in generally known textbooks and handbooks. The
textbook by Winnacker: Gene und Klone, Eine Einführung in die
Gentechnologie (Verlag Chemie, Weinheim, Germany, 1990), or
the handbook by Sambrook et al.: Molecular Cloning, A
Laboratory Manual (Cold Spring Harbor Laboratory Press, 1989)
may be mentioned as example. A well-known gene library is that
of the E. coli K-12 strain W3110 set up in λ vectors by Kohara
et al. (Cell 50, 495 -508 (1987)). Bathe et al. (Molecular and
25 General Genetics, 252:255-265, 1996) describe a gene library
of C. glutamicum ATCC13032, which was set up with the aid of
the cosmid vector SuperCos I (Wahl et al., 1987, Proceedings
of the National Academy of Sciences USA, 84:2160-2164) in the
E. coli K-12 strain NM554 (Raleigh et al., 1988, Nucleic Acids
30 Research 16:1563-1575).

Börmann et al. (Molecular Microbiology 6(3), 317-326) (1992))
in turn describe a gene library of C. glutamicum ATCC13032
using the cosmid pHc79 (Hohn and Collins, Gene 11, 291-298
(1980)). To prepare a gene library of C. glutamicum in E. coli
35 it is also possible to use plasmids such as pBR322 (Bolivar,

Life Sciences, 25, 807-818 (1979)) or pUC9 (Vieira et al., 1982, Gene, 19:259-268). Suitable hosts are, in particular, those E. coli strains which are restriction- and recombination-defective. An example of these is the strain
5 DH5 α mcr, which has been described by Grant et al. (Proceedings of the National Academy of Sciences USA, 87 (1990) 4645-4649). The long DNA fragments cloned with the aid of cosmids can in turn be subcloned in the usual vectors suitable for sequencing and then sequenced, as is described e.g. by Sanger et al.
10 (Proceedings of the National Academy of Sciences of the United States of America, 74:5463-5467, 1977).

The resulting DNA sequences can then be investigated with known algorithms or sequence analysis programs, such as that of Staden (Nucleic Acids Research 14, 217-232(1986)), that of
15 Marck (Nucleic Acids Research 16, 1829-1836 (1988)) or the GCG program of Butler (Methods of Biochemical Analysis 39, 74-97 (1998)).

The new DNA sequence of C. glutamicum which codes for the metE gene and which, as SEQ ID No. 1, is a constituent of the
20 present invention has been found. The amino acid sequence of the corresponding protein has furthermore been derived from the present DNA sequence by the methods described above. The resulting amino acid sequence of the metE gene product is shown in SEQ ID No. 2.

25 Coding DNA sequences which result from SEQ ID No. 1 by the degeneracy of the genetic code are also a constituent of the invention. In the same way, DNA sequences which hybridize with SEQ ID No. 1 or parts of SEQ ID No. 1 are a constituent of the invention. Conservative amino acid exchanges, such as e.g.
30 exchange of glycine for alanine or of aspartic acid for glutamic acid in proteins, are furthermore known among experts as "sense mutations" which do not lead to a fundamental change in the activity of the protein, i.e. they are of neutral function.

35 It is furthermore known that changes at the N and/or C terminus of a protein must not substantially impair and may

even stabilize the function thereof. Information in this context can be found in Ben-Bassat et al. (Journal of Bacteriology 169:751-757 (1987)), in O'Regan et al. (Gene 77:237-251 (1989)), in Sahin-Toth et al. (Protein Sciences 3:240-247 (1994)), in Hochuli et al. (Bio/Technology 6:1321-1325 (1988)) and in known textbooks of genetics and molecular biology. Amino acid sequences which result in a corresponding manner from SEQ ID No. 2 are also a constituent of the invention.

In the same way, DNA sequences which hybridize with SEQ ID No. 1 or parts of SEQ ID No. 1 are a constituent of the invention. Finally, DNA sequences which are prepared by the polymerase chain reaction (PCR) using primers which result from SEQ ID No. 1 are a constituent of the invention. Such oligonucleotides typically have a length of at least 15 nucleotides.

Instructions for identifying DNA sequences by means of hybridization can be found in the handbook "The DIG System Users Guide for Filter Hybridization" from Boehringer Mannheim GmbH (Mannheim, Germany, 1993) and in Liebl et al. (International Journal of Systematic Bacteriology (1991) 41: 255-260). Instructions for amplification of DNA sequences with the aid of the polymerase chain reaction (PCR) can be found in the handbook by Gait: Oligonucleotide synthesis: A Practical Approach (IRL Press, Oxford, UK, 1984) and in Newton and Graham: PCR (Spektrum Akademischer Verlag, Heidelberg, Germany, 1994).

It has been found that coryneform bacteria produce amino acids, in particular L-methionine, in an improved manner after over-expression of the metE gene.

To achieve an over-expression, the number of copies of the corresponding genes can be increased, or the promoter and regulation region or the ribosome binding site upstream of the structural gene can be mutated. Expression cassettes which are incorporated upstream of the structural gene act in the same way. By inducible promoters, it is additionally possible to

increase the expression in the course of fermentative L-methionine production. The expression is likewise improved by measures to prolong the life of the m-RNA. Furthermore, the enzyme activity is also increased by preventing the degradation of the enzyme protein. The genes or gene constructs can either be present in plasmids with a varying number of copies, or can be integrated and amplified in the chromosome. Alternatively, an over-expression of the genes in question can furthermore be achieved by changing the composition of the media and the culture procedure.

Instructions in this context can be found in Martin et al. (Bio/Technology 5, 137-146 (1987)), in Guerrero et al. (Gene 138, 35-41 (1994)), Tsuchiya and Morinaga (Bio/Technology 6, 428-430 (1988)), in Eikmanns et al. (Gene 102, 93-98 (1991)), in European Patent Specification 0 472 869, in US Patent 4,601,893, in Schwarzer and Pühler (Bio/Technology 9, 84-87 (1991)), in Reinscheid et al. (Applied and Environmental Microbiology 60, 126-132 (1994)), in LaBarre et al. (Journal of Bacteriology 175, 1001-1007 (1993)), in Patent Application WO 96/15246, in Malumbres et al. (Gene 134, 15 - 24 (1993)), in Japanese Laid-Open Specification JP-A-10-229891, in Jensen and Hammer (Biotechnology and Bioengineering 58, 191-195 (1998)), in Makrides (Microbiological Reviews 60:512-538 (1996)) and in known textbooks of genetics and molecular biology.

By way of example, for enhancement the metE gene according to the invention was over-expressed with the aid of episomal plasmids. Suitable plasmids are those which are replicated in coryneform bacteria. Numerous known plasmid vectors, such as e.g. pZ1 (Menkel et al., Applied and Environmental Microbiology (1989) 64: 549-554), pEKEx1 (Eikmanns et al., Gene 102:93-98 (1991)) or pHS2-1 (Sonnen et al., Gene 107:69-74 (1991)) are based on the cryptic plasmids pHM1519, pBL1 or pGA1. Other plasmid vectors, such as those based on pCG4 (US-A 4,489,160), or pNG2 (Serwold-Davis et al., FEMS Microbiology Letters 66, 119-124 (1990)), or pAG1 (US-A 5,158,891), can be used in the same manner.

Plasmid vectors which are furthermore suitable are also those with the aid of which the process of gene amplification by integration into the chromosome can be used, as has been described, for example, by Reinscheid et al. (Applied and Environmental Microbiology 60, 126-132 (1994)) for duplication or amplification of the hom-thrB operon. In this method, the complete gene is cloned in a plasmid vector which can replicate in a host (typically E. coli), but not in C. glutamicum. Possible vectors are, for example, pSUP301 (Simon et al., Bio/Technology 1, 784-791 (1983)), pK18mob or pK19mob (Schäfer et al., Gene 145, 69-73 (1994)), pGEM-T (Promega corporation, Madison, WI, USA), pCR2.1-TOPO (Shuman (1994). Journal of Biological Chemistry 269:32678-84; US-A 5,487,993), pCR®Blunt (Invitrogen, Groningen, Holland; Bernard et al., Journal of Molecular Biology, 234: 534-541 (1993)), pEM1 (Schrumpf et al, 1991, Journal of Bacteriology 173:4510-4516) or pBGS8 (Spratt et al., 1986, Gene 41: 337-342). The plasmid vector which contains the gene to be amplified is then transferred into the desired strain of C. glutamicum by conjugation or transformation. The method of conjugation is described, for example, by Schäfer et al. (Applied and Environmental Microbiology 60, 756-759 (1994)). Methods for transformation are described, for example, by Thierbach et al. (Applied Microbiology and Biotechnology 29, 356-362 (1988)), Dunican and Shivnan (Bio/Technology 7, 1067-1070 (1989)) and Tauch et al. (FEMS Microbiological Letters 123, 343-347 (1994)). After homologous recombination by means of a "cross over" event, the resulting strain contains at least two copies of the gene in question.

In addition, it may be advantageous for the production of amino acids, in particular L-methionine, to enhance one or more enzymes of the particular biosynthesis pathway, of glycolysis, of anaplerosis, of the citric acid cycle or of amino acid export, in addition to the metE gene.

Thus for the preparation of amino acids, in particular L-methionine, one or more genes chosen from the group consisting of

10
15
20

- the gap gene which codes for glyceraldehyde 3-phosphate dehydrogenase (Eikmanns (1992), Journal of Bacteriology 174:6076-6086),
- the tpi gene which codes for triose phosphate isomerase (Eikmanns (1992), Journal of Bacteriology 174:6076-6086),
- the pgk gene which codes for 3-phosphoglycerate kinase (Eikmanns (1992), Journal of Bacteriology 174:6076-6086),
- the pyc gene which codes for pyruvate carboxylase (Eikmanns (1992), Journal of Bacteriology 174:6076-6086),
- the lysC gene which codes for a feed-back resistant aspartate kinase (Accession No.P26512; EP-B-0387527; EP-A-0699759),
- the metaA gene which codes for homoserine O-acetyltransferase (ACCESSION Number AF052652),
- the metB gene which codes for cystathionine gamma-synthase (ACCESSION Number AF126953),
- the aecD gene which codes for cystathionine gamma-lyase (ACCESSION Number M89931)
- the glyA gene which codes for serine hydroxymethyltransferase (JP-A-08107788),
- the metY gene which codes for O-acetylhomoserine sulphydrylase (DSM 13556)

can be enhanced, in particular over-expressed.

It may furthermore be advantageous for the production of amino acids, in particular L-methionine, in addition to the enhancement of the metE gene, for one or more genes chosen from the group consisting of

- the thrB gene which codes for homoserine kinase (ACCESSION Number P08210),

- the *ilvA* gene which codes for threonine dehydratase (ACCESSION Number Q04513),
- the *thrC* gene which codes for threonine synthase (ACCESSION Number P23669),
- the *ddh* gene which codes for meso-diaminopimelate D-dehydrogenase (ACCESSION Number Y00151),
- the *pck* gene which codes for phosphoenol pyruvate carboxykinase (DE 199 50 409.1; DSM 13047),
- the *pgi* gene which codes for glucose 6-phosphate isomerase (US 09/396,478; DSM 12969),
- the *poxB* gene which codes for pyruvate oxidase (DE: 1995 1975.7; DSM 13114)

to be attenuated, in particular for the expression thereof to be reduced.

The term "attenuation" in this connection describes the reduction or elimination of the intracellular activity of one or more enzymes (proteins) in a microorganism which are coded by the corresponding DNA, for example by using a weak promoter or using a gene or allele which codes for a corresponding enzyme with a low activity or inactivates the corresponding gene or enzyme (protein), and optionally combining these measures.

By attenuation measures, the activity or concentration of the corresponding protein is in general reduced to 0 to 50%, 0 to 25%, 0 to 10% or 0 to 5% of the activity or concentration of the wild-type protein.

In addition to over-expression of the metE gene it may furthermore be advantageous for the production of amino acids, in particular L-methionine, to eliminate undesirable side reactions, (Nakayama: "Breeding of Amino Acid Producing Microorganisms", in: Overproduction of Microbial Products,

Krumphanzl, Sikyta, Vanek (eds.), Academic Press, London, UK, 1982).

5 The microorganisms prepared according to the invention can be cultured continuously or discontinuously in the batch process (batch culture) or in the fed batch (feed process) or repeated fed batch process (repetitive feed process) for the purpose of production of amino acids, in particular L-methionine. A summary of known culture methods is described in the textbook by Chmiel (Bioprozesstechnik 1. Einführung in die Bioverfahrenstechnik (Gustav Fischer Verlag, Stuttgart, 1991)) or in the textbook by Storhas (Bioreaktoren und periphere Einrichtungen (Vieweg Verlag, Braunschweig/Wiesbaden, 1994)).

10 The culture medium to be used must meet the requirements of the particular strains in a suitable manner. Descriptions of culture media for various microorganisms are contained in the handbook "Manual of Methods for General Bacteriology" of the American Society for Bacteriology (Washington D.C., USA, 1981).

15 Sugars and carbohydrates, such as e.g. glucose, sucrose, lactose, fructose, maltose, molasses, starch and cellulose, oils and fats, such as e.g. soya oil, sunflower oil, groundnut oil and coconut fat, fatty acids, such as e.g. palmitic acid, stearic acid and linoleic acid, alcohols, such as e.g. glycerol and ethanol, and organic acids, such as e.g. acetic acid, can be used as the source of carbon. These substance can be used individually or as a mixture.

20 Organic nitrogen-containing compounds, such as peptones, yeast extract, meat extract, malt extract, corn steep liquor, soya bean flour and urea, or inorganic compounds, such as ammonium sulfate, ammonium chloride, ammonium phosphate, ammonium carbonate and ammonium nitrate, can be used as the source of nitrogen. The sources of nitrogen can be used individually or as a mixture.

25 Organic and inorganic sulfur-containing compounds, such as, for example, sulfides, sulfites, sulfates and thiosulfates,

can be used as a source of sulfur, in particular for the preparation of methionine.

Phosphoric acid, potassium dihydrogen phosphate or dipotassium hydrogen phosphate or the corresponding sodium-containing salts can be used as the source of phosphorus. The culture medium must furthermore comprise salts of metals, such as e. g. magnesium sulfate or iron sulfate, which are necessary for growth. Finally, essential growth substances, such as amino acids and vitamins, can be employed in addition to the above-mentioned substances. Suitable precursors can moreover be added to the culture medium. The starting substances mentioned can be added to the culture in the form of a single batch, or can be fed in during the culture in a suitable manner.

Basic compounds, such as sodium hydroxide, potassium hydroxide, ammonia or aqueous ammonia, or acid compounds, such as phosphoric acid or sulfuric acid, can be employed in a suitable manner to control the pH of the culture. Antifoams, such as e.g. fatty acid polyglycol esters, can be employed to control the development of foam. Suitable substances having a selective action, such as e.g. antibiotics, can be added to the medium to maintain the stability of plasmids. To maintain aerobic conditions, oxygen or oxygen-containing gas mixtures, such as e.g. air, are introduced into the culture. The temperature of the culture is usually 20°C to 45°C, and preferably 25°C to 40°C. Culturing is continued until a maximum of the desired product has formed. This target is usually reached within 10 hours to 160 hours.

The fermentation broths obtained in this way, in particular containing L-methionine, usually have a dry weight of 7.5 to 25 wt.% and contain L-methionine. It is furthermore also advantageous if the fermentation is conducted in a sugar-limited procedure at least at the end, but in particular over at least 30% of the duration of the fermentation. That is to say, the concentration of utilizable sugar in the fermentation medium is reduced to ≥ 0 to 3 g/l during this period.

The fermentation broth prepared in this manner, in particular containing L-methionine, is then further processed. Depending on requirements all or some of the biomass can be removed from the fermentation broth by separation methods, such as

This free-flowing, finely divided powder can then in turn be converted by suitable compacting or granulating processes into a coarse-grained, readily free-flowing, storable and largely dust-free product. In the granulation or compacting it is advantageous to employ conventional organic or inorganic auxiliary substances or carriers, such as starch, gelatin, cellulose derivatives or similar substances, such as are conventionally used as binders, gelling agents or thickeners in foodstuffs or feedstuffs processing, or further substances, such as, for example, silicas, silicates or stearates.

"Free-flowing" is understood as meaning powders which flow unimpeded out of the vessel with the opening of 5 mm (millimeters) of a series of glass outflow vessels with outflow openings of various sizes (Klein, Seifen, Öle, Fette, Wachse 94, 12 (1968)).

As described here, "finely divided" means a powder with a predominant content (> 50 %) having a particle size of 20 to 200 µm diameter. "Coarse-grained" means products with a predominant content (> 50 %) having a particle size of 200 to 2000 µm diameter. In this context, "dust-free" means that the product contains only small contents (< 5 %) having particle sizes of less than 20 µm diameter. The particle size determination can be carried out with methods of laser

preferably up to 100%. In another aspect of the invention, up to 20% of the biomass, preferably up to 15%, preferably up to 10%, preferably up to 5%, particularly preferably no biomass is separated off.

- 5 These organic substances include organic by-products which are optionally produced, in addition to the L-methionine, and optionally discharged by the microorganisms employed in the fermentation. These include L-amino acids chosen from the group consisting of L-lysine, L-valine, L-threonine, L-alanine
10 or L-tryptophan. They include vitamins chosen from the group consisting of vitamin B1 (thiamine), vitamin B2 (riboflavin), vitamin B5 (pantothenic acid), vitamin B6 (pyridoxine), vitamin B12 (cyanocobalamin), nicotinic acid/nicotinamide and vitamin E (tocopherol). They also
15 include organic acids which carry one to three carboxyl groups, such as, acetic acid, lactic acid, citric acid, malic acid or fumaric acid. Finally, they also include sugars, such as, for example, trehalose. These compounds are optionally desired if they improve the nutritional value of the product.
- 20 These organic substances, including L-methionine and/or D-methionine and/or the racemic mixture D,L-methionine, can also be added, depending on requirements, as a concentrate or pure substance in solid or liquid form during a suitable process step. These organic substances mentioned can be added
25 individually or as mixtures to the resulting or concentrated fermentation broth, or also during the drying or granulation process. It is likewise possible to add an organic substance or a mixture of several organic substances to the fermentation broth and a further organic substance or a further mixture of
30 several organic substances during a later process step, for example granulation.

The product described above is suitable as a feedstuffs additive, i.e. feed additive, for animal nutrition.

- 35 The L-methionine content of the animal feedstuffs additive is conventionally 1 wt.% to 80 wt.%, preferably 2 wt.% to 80 wt.%, particularly preferably 4 wt.% to 80 wt.%, and very

particularly preferably 8 wt.% to 80 wt.%, based on the dry weight of the animal feedstuffs additive. Contents of 1 wt.% to 60 wt.%, 2 wt.% to 60 wt.%, 4 wt.% to 60 wt.%, 6 wt.% to 60 wt.%, 1 wt.% to 40 wt.%, 2 wt.% to 40 wt.% or 4 wt.% to 40 wt.% are likewise possible. The water content of the feedstuffs additive is conventionally up to 5 wt.%, preferably up to 4 wt.%, and particularly preferably less than 2 wt.%.
5

The invention also provides a process for the preparation of an L-methionine-containing animal feedstuffs additive from fermentation broths, which comprises the steps
10

- a) culture and fermentation of an L-methionine-producing microorganism in a fermentation medium;
- b) removal of water from the L-methionine-containing fermentation broth (concentration);
- 15 c) removal of an amount of 0 to 100 wt.% of the biomass formed during the fermentation; and
- d) drying of the fermentation broth obtained according to a) and/or b) to obtain the animal feedstuffs additive in the desired powder or granule form.

20 If desired, one or more of the following steps can furthermore be carried out in the process according to the invention:

- 25 e) addition of one or more organic substances, including L-methionine and/or D-methionine and/or the racemic mixture D,L-methionine, to the products obtained according to a), b) and/or c);
- f) addition of auxiliary substances chosen from the group consisting of silicas, silicates, stearates, grits and bran to the substances obtained according to a) to d) for stabilization and to increase the storability; or
- 30 g) conversion of the substances obtained according to a) to e) into a form stable to the animal stomach, in particular rumen, by coating with film-forming agents.

The analysis of L-methionine can be carried out by ion exchange chromatography with subsequent ninhydrin derivation, as described by Spackman et al. (Analytical Chemistry, 30, (1958), 1190).

- 5 The process according to the invention is used for the fermentative preparation of amino acids, in particular L-methionine.

10 The following microorganisms were deposited as a pure culture on 14th June 2001 at the Deutsche Sammlung für Mikroorganismen und Zellkulturen (DSMZ = German Collection of Microorganisms and Cell Cultures, Braunschweig, Germany) in accordance with the Budapest Treaty:

- Escherichia coli DH5 α mcr/pCREmetAE as DSM 14352,
- Escherichia coli DH5 α mcr/pCREmetAEY as DSM 14353.

15 The present invention is explained in more detail in the following with the aid of embodiment examples.

Example 1

Preparation of a genomic cosmid gene library from Corynebacterium glutamicum ATCC 13032

20 Chromosomal DNA from Corynebacterium glutamicum ATCC-13032 was isolated as described by Tauch et al. (1995, Plasmid 33:168-179) and partly cleaved with the restriction enzyme Sau3AI (Amersham Pharmacia, Freiburg, Germany, Product Description Sau3AI, Code no. 27-0913-02). The DNA fragments were
25 dephosphorylated with shrimp alkaline phosphatase (Roche Diagnostics GmbH, Mannheim, Germany, Product Description SAP, Code no. 1758250). The DNA of the cosmid vector SuperCos1 (Wahl et al. (1987) Proceedings of the National Academy of Sciences USA 84:2160-2164), obtained from Stratagene (La
30 Jolla, USA, Product Description SuperCos1 Cosmid Vector Kit, Code no. 251301) was cleaved with the restriction enzyme XbaI (Amersham Pharmacia, Freiburg, Germany, Product Description

XbaI, Code no. 27-0948-02) and likewise dephosphorylated with shrimp alkaline phosphatase.

The cosmid DNA was then cleaved with the restriction enzyme BamHI (Amersham Pharmacia, Freiburg, Germany, Product Description BamHI, Code no. 27-0868-04). The cosmid DNA treated in this manner was mixed with the treated ATCC13032 DNA and the batch was treated with T4 DNA ligase (Amersham Pharmacia, Freiburg, Germany, Product Description T4-DNA-Ligase, Code no.27-0870-04). The ligation mixture was then packed in phages with the aid of Gigapack II XL Packing Extract (Stratagene, La Jolla, USA, Product Description Gigapack II XL Packing Extract, Code no. 200217).

For infection of the E. coli strain NM554 (Raleigh et al. 1988, Nucleic Acid Research 16:1563-1575) the cells were taken up in 10 mM MgSO₄ and mixed with an aliquot of the phage suspension. The infection and titering of the cosmid library were carried out as described by Sambrook et al. (1989, Molecular Cloning: A laboratory Manual, Cold Spring Harbor), the cells being plated out on LB agar (Lennox, 1955, Virology, 1:190) with 100 mg/l ampicillin. After incubation overnight at 37°C, recombinant individual clones were selected.

Example 2

Isolation and sequencing of the metE gene

The cosmid DNA of an individual colony was isolated with the Qiaprep Spin Miniprep Kit (Product No. 27106, Qiagen, Hilden, Germany) in accordance with the manufacturer's instructions and partly cleaved with the restriction enzyme Sau3AI (Amersham Pharmacia, Freiburg, Germany, Product Description Sau3AI, Product No. 27-0913-02). The DNA fragments were dephosphorylated with shrimp alkaline phosphatase (Roche Diagnostics GmbH, Mannheim, Germany, Product Description SAP, Product No. 1758250). After separation by gel electrophoresis, the cosmid fragments in the size range of 1500 to 2000 bp were isolated with the QiaExII Gel Extraction Kit (Product No. 20021, Qiagen, Hilden, Germany).

0901995 "1995" 1995

The DNA of the sequencing vector pZero-1, obtained from Invitrogen (Groningen, The Netherlands, Product Description Zero Background Cloning Kit, Product No. K2500-01) was cleaved with the restriction enzyme BamHI (Amersham Pharmacia, Freiburg, Germany, Product Description BamHI, Product No. 27-0868-04). The ligation of the cosmid fragments in the sequencing vector pZero-1 was carried out as described by Sambrook et al. (1989, Molecular Cloning: A Laboratory Manual, Cold Spring Harbor), the DNA mixture being incubated overnight with T4 ligase (Pharmacia Biotech, Freiburg, Germany). This ligation mixture was then electroporated (Tauch et al. 1994, FEMS Microbiol Letters, 123:343-7) into the E. coli strain DH5 α mc^r (Grant, 1990, Proceedings of the National Academy of Sciences U.S.A., 87:4645-4649) and plated out on LB agar (Lennox, 1955, Virology, 1:190) with 50 mg/l zeocin.

The plasmid preparation of the recombinant clones was carried out with Biorobot 9600 (Product No. 900200, Qiagen, Hilden, Germany). The sequencing was carried out by the dideoxy chain termination method of Sanger et al. (1977, Proceedings of the National Academy of Sciences U.S.A., 74:5463-5467) with modifications according to Zimmermann et al. (1990, Nucleic Acids Research, 18:1067). The "RR dRhodamin Terminator Cycle Sequencing Kit" from PE Applied Biosystems (Product No. 403044, Weiterstadt, Germany) was used. The separation by gel electrophoresis and analysis of the sequencing reaction were carried out in a "Rotiphoresis NF Acrylamide/Bisacrylamide" Gel (29:1) (Product No. A124.1, Roth, Karlsruhe, Germany) with the "ABI Prism 377" sequencer from PE Applied Biosystems (Weiterstadt, Germany).

The raw sequence data obtained were then processed using the Staden program package (1986, Nucleic Acids Research, 14:217-231) version 97-0. The individual sequences of the pZero1 derivatives were assembled to a continuous contig. The computer-assisted coding region analysis was prepared with the XNIP program (Staden, 1986, Nucleic Acids Research, 14:217-231).

The resulting nucleotide sequence is shown in SEQ ID No. 1. Analysis of the nucleotide sequence showed an open reading frame of 2235 base pairs, which was called the metE gene. The metE gene codes for a protein of 745 amino acids.

5 Example 3

Preparation of the strains *C. glutamicum* ATCC13032/pCREmetA and ATCC13032/pCREmetAE

3.1 Amplification of the genes metA and metE

From the strain ATCC13032, chromosomal DNA was isolated by the method of Eikmanns et al. (Microbiology 140: 1817 -1828 (1994)). Starting from the nucleotide sequences of the methionine biosynthesis genes metA (gene library Accession Number AF052652) and metE (SEQ ID No. 1) of *C. glutamicum* ATCC13032, the following oligonucleotides were chosen for the polymerase chain reaction (PCR) (see SEQ ID No. 3, SEQ ID No. 4, SEQ ID No. 5 and SEQ ID No. 6):

metA-EVP5:

5'- AGAACGAATTCAAAGGAGGACAACCATGCCCCACCCTCGCGC -3'

metA-EVP3:

5'- GTCGTGGATCCCCTATTAGATGTAGAACTCG -3'

metE-EVP5:

5'-GGCTCAAAGATCTAAAGGAGGACAACCATGACTTCCAACCTTTCTTC -3'

metE-EVP3:

5'- GGTTCCCTGTCGACGGTACCATTTAGATAGTTGCTCCGATT -3'

25 The primers shown were synthesized by MWG-Biotech AG (Ebersberg, Germany) and the PCR reaction was carried out by the standard PCR method of Innis et al. (PCR Protocols. A Guide to Methods and Applications, 1990, Academic Press) with Pwo-Polymerase from Roche Diagnostics GmbH (Mannheim, Germany). With the aid of the polymerase chain reaction, the
30 primers allow amplification of a DNA fragment 1161 bp in size,

which carries the metA gene, and a DNA fragment 2286 bp in size, which carries the metE gene.

Furthermore, the primer metA-EVP5 contains the sequence for the cleavage site of the restriction endonuclease EcoRI, the primer metA-EVP3 the cleavage site of the restriction endonuclease BamHI, the primer metE-EVP5 the cleavage site of the restriction endonuclease BglII and the primer metE-EVP3 the cleavage site of the restriction endonuclease SalI, which are marked by underlining in the nucleotide sequence shown above.

The metA fragment 1161 bp in size was cleaved with the restriction endonucleases EcoRI and BamHI, and the metE fragment 2286 bp in size was cleaved with the restriction endonucleases BglII and SalI. The two batches were separated by gel electrophoresis and the fragments metA (approx. 1150 bp) and metE (approx. 2270 bp) were then isolated from the agarose gel with the QiaExII Gel Extraction Kit (Product No. 20021, Qiagen, Hilden, Germany).

3.2 Cloning of metA in the vector pZ8-1

The E. coli - C. glutamicum shuttle expression vector pZ8-1 (EP 0 375 889) was used as the base vector for the expression.

DNA of the plasmid pZ8-1 was cleaved completely with the restriction enzymes EcoRI and BamHI and then dephosphorylated with shrimp alkaline phosphatase (Roche Diagnostics GmbH, Mannheim, Germany, Product Description SAP, Product No. 1758250).

The metA fragment approx. 1150 bp in size isolated from the agarose gel in example 3.1 and cleaved with the restriction endonucleases BamHI and EcoRI was mixed with the vector pZ8-1 prepared in this way and the batch was treated with T4 DNA ligase (Amersham Pharmacia, Freiburg, Germany, Product Description T4-DNA-Ligase, Code no.27-0870-04).

The ligation batch was transformed in the E. coli strain DH5 α mcr (Hanahan, In: DNA cloning. A Practical Approach. Vol.

I. IRL-Press, Oxford, Washington DC, USA). Selection of plasmid-carrying cells was made by plating out the transformation batch on LB agar (Lennox, 1955, Virology, 1:190) with 50 mg/l kanamycin. After incubation overnight at 37°C, recombinant individual clones were selected. Plasmid DNA was isolated from a transformant with the Qiaprep Spin Miniprep Kit (Product No. 27106, Qiagen, Hilden, Germany) in accordance with the manufacturer's instructions and checked by restriction cleavage. The resulting plasmid was called pCREmetA.

3.3 Cloning of metE in the vector pCREmetA

The plasmid pCREmetA described in example 3.2 was cleaved completely with the restriction enzymes BamHI and SalI and then dephosphorylated with shrimp alkaline phosphatase (Roche Diagnostics GmbH, Mannheim, Germany, Product Description SAP, Product No. 1758250).

The metE fragment approx. 2270 bp in size obtained in example 3.1 by means of the polymerase chain reaction and cleaved with the restriction endonucleases BglII and SalI was mixed with the vector pCREmetA prepared in this way. The batch was treated with T4 DNA ligase (Amersham Pharmacia, Freiburg, Germany, Product Description T4-DNA-Ligase, Code no.27-0870-04).

The ligation batch was transformed in the E. coli strain DH5 α mc^r (Hanahan, In: DNA cloning. A Practical Approach. Vol. I. IRL-Press, Oxford, Washington DC, USA). Selection of plasmid-carrying cells was made by plating out the transformation batch on LB agar (Lennox, 1955, Virology, 1:190) with 50 mg/l kanamycin. After incubation overnight at 37°C, recombinant individual clones were selected. Plasmid DNA was isolated from a transformant with the Qiaprep Spin Miniprep Kit (Product No. 27106, Qiagen, Hilden, Germany) in accordance with the manufacturer's instructions and checked by restriction cleavage. The resulting plasmid was called pCREmetAE. It is shown in figure 1. The strain E. coli DH5 α mc^r/pCREmetAE was deposited as a pure culture on 14th June

2001 at the Deutsche Sammlung für Mikroorganismen und Zellkulturen (DSMZ = German Collection of Microorganisms and Cell Cultures, Braunschweig, Germany) in accordance with the Budapest Treaty as DSM 14352.

5 3.4 Preparation of the strains *C. glutamicum*
ATCC23032/pCREmetA and ATCC13032/pCREmetAE

The vectors pCREmetA and pCREmetAE obtained in example 3.2 and 3.3 were electroporated in the strain *C. glutamicum* ATCC13032 using the electroporation method described by Liebl et al.

10 (FEMS Microbiology Letters, 53:299-303 (1989)). Selection of the plasmid-carrying cells took place on LBHIS agar comprising 18.5 g/l brain-heart infusion broth, 0.5 M sorbitol, 5 g/l Bacto-tryptone, 2.5 g/l Bacto-yeast extract, 5 g/l NaCl and 18 g/l Bacto-agar, which had been supplemented with 25 mg/l kanamycin. Incubation was carried out for 2 days at 33°C.

15 Plasmid DNA was isolated from in each case one transformant by conventional methods (Peters-Wendisch et al., 1998, Microbiology 144, 915-927) and checked by restriction cleavage. The resulting strains were called ATCC13032/pCREmetA and ATCC13032/pCREmetAE.

20 Example 4

Preparation of L-methionine with the strain *C. glutamicum* ATCC13032/pCREmetAE

25 The *C. glutamicum* strains ATCC13032/pCREmetA and ATCC13032/pCREmetAE obtained in example 3 were cultured in a nutrient medium suitable for the production of methionine and the methionine content in the culture supernatant was determined.

30 For this, the strains were first incubated on an agar plate with the corresponding antibiotic (brain-heart agar with kanamycin (50 mg/l)) for 24 hours at 33°C. Starting from this agar plate culture, in each case a preculture was seeded (10 ml medium in a 100 ml conical flask). The complete medium CgIII was used as the medium for the precultures.

Medium Cg III

NaCl	2.5 g/l
Bacto-Peptone	10 g/l
Bacto-Yeast extract	10 g/l
Glucose (autoclaved separately)	2% (w/v)

The pH was brought to pH 7.4

Kanamycin (25 mg/l) was added to this. The preculture was incubated for 16 hours at 33°C at 240 rpm on a shaking machine. In each case a main culture was seeded from these precultures such that the initial OD (660 nm) of the main cultures was 0.1. Medium MM was used for the main cultures.

Medium MM

CSL (corn steep liquor)	5 g/l
MOPS (morpholinopropanesulfonic acid)	20 g/l
Glucose (autoclaved separately)	50 g/l
(NH ₄) ₂ SO ₄	25 g/l
KH ₂ PO ₄	0.1 g/l
MgSO ₄ * 7 H ₂ O	1.0 g/l
CaCl ₂ * 2 H ₂ O	10 mg/l
FeSO ₄ * 7 H ₂ O	10 mg/l
MnSO ₄ * H ₂ O	5.0mg/l
Biotin (sterile-filtered)	0.3 mg/l
Thiamine * HCl (sterile-filtered)	0.2 mg/l
CaCO ₃	25 g/l

The CSL, MOPS and the salt solution were brought to pH 7 with aqueous ammonia and autoclaved. The sterile substrate and vitamin solutions were then added, as well as the CaCO₃ autoclaved in the dry state.

Culturing is carried out in a 10 ml volume in 100 ml conical flasks with baffles. Kanamycin (25 mg/l) was added. Culturing was carried out at 33°C and 80% atmospheric humidity.

After 72 hours, the OD was determined at a measurement wavelength of 660 nm with a Biomek 1000 (Beckmann Instruments GmbH, Munich). The amount of methionine formed was determined with an amino acid analyzer from Eppendorf-BioTronik (Hamburg, Germany) by ion exchange chromatography and post-column derivation with ninhydrin detection.

The result of the experiment is shown in Table 1.

Table 1

Strain	OD (660 nm)	Methionine mg/l
ATCC13032/pCREmetA	12.3	6.6
ATCC13032/pCREmetAE	14.3	15.3

Example 5

Preparation of the strain *C. glutamicum* ATCC13032/pCREmetAEY

5.1 Amplification of the metY gene

From the strain ATCC13032, chromosomal DNA was isolated by the method of Eikmanns et al. (Microbiology 140: 1817 -1828 (1994)). Starting from the nucleotide sequence of the methionine biosynthesis gene metY (DE: 10043334.0) of *C. glutamicum* ATCC13032, the following oligonucleotides were chosen for the polymerase chain reaction (PCR) (see SEQ ID No. 7 and SEQ ID No. 8):

metY-EVP5:

5'- CTAATAAGTCGACAAAGGAGGACAACCATGCCAAAGTACGAC -3'

metY-EVP3:

5'- GAGTCTAATGCATGCTAGATTGCAGCAAAGCCG -3'

The primers shown were synthesized by MWG-Biotech AG (Ebersberg, Germany) and the PCR reaction was carried out by the standard PCR method of Innis et al. (PCR Protocols. A Guide to Methods and Applications, 1990, Academic Press) with Pwo-Polymerase from Roche Diagnostics GmbH (Mannheim, Germany). With the aid of the polymerase chain reaction, the primers allow amplification of a DNA fragment 1341 bp in size, which carries the metY gene.

Furthermore, the primer metY-EVP5 contains the sequence for the cleavage site of the restriction endonuclease SalI and the

primer metY-EVP3 the cleavage site of the restriction endonuclease NsiI, which are marked by underlining in the nucleotide sequence shown above.

5 The metY fragment 1341 bp in size was cleaved with the restriction endonucleases SalI and NsiI. The batch was separated by gel electrophoresis and the fragment metY (approx. 1330 bp) was then isolated from the agarose gel with the QiaExII Gel Extraction Kit (Product No. 20021, Qiagen, Hilden, Germany).

10 5.2 Cloning of metA and metY in the vector pZ8-1

The plasmid pCREmetA described in example 3.2 was cleaved completely with the restriction enzymes SalI and PstI and then dephosphorylated with shrimp alkaline phosphatase (Roche Diagnostics GmbH, Mannheim, Germany, Product Description SAP, Product No. 1758250).

The metY fragment approx. 1330 bp in size isolated from the agarose gel in example 5.1 and cleaved with the restriction endonucleases SalI and NsiI was mixed with the vector pCREmetA prepared in this way and the batch was treated with T4 DNA ligase (Amersham Pharmacia, Freiburg, Germany, Product Description T4-DNA-Ligase, Code no.27-0870-04).

25 The ligation batch was transformed in the E. coli strain DH5 α mc^r (Hanahan, In: DNA cloning. A Practical Approach. Vol. I. IRL-Press, Oxford, Washington DC, USA). Selection of plasmid-carrying cells was made by plating out the transformation batch on LB agar (Lennox, 1955, Virology, 1:190) with 50 mg/l kanamycin. After incubation overnight at 37°C, recombinant individual clones were selected. Plasmid DNA was isolated from a transformant with the Qiaprep Spin
30 Miniprep Kit (Product No. 27106, Qiagen, Hilden, Germany) in accordance with the manufacturer's instructions and checked by restriction cleavage. The resulting plasmid was called pCREmetAY.

5.3 Cloning of metE in the vector pCREmetAY

The plasmid pCREmetAY described in example 5.2 was cleaved completely with the restriction enzymes BamHI and SalI and then dephosphorylated with shrimp alkaline phosphatase (Roche Diagnostics GmbH, Mannheim, Germany, Product Description SAP, Product No. 1758250).

The metE fragment approx. 2270 bp in size obtained in example 3.1 by means of the polymerase chain reaction and cleaved with the restriction endonucleases BglII and SalI was mixed with the vector pCREmetAY prepared in this way. The batch was treated with T4 DNA ligase (Amersham Pharmacia, Freiburg, Germany, Product Description T4-DNA-Ligase, Code no.27-0870-04).

The ligation batch was transformed in the E. coli strain DH5 α mcrr (Hanahan, In: DNA cloning. A Practical Approach. Vol. I. IRL-Press, Oxford, Washington DC, USA). Selection of plasmid-carrying cells was made by plating out the transformation batch on LB agar (Lennox, 1955, Virology, 1:190) with 50 mg/l kanamycin. After incubation overnight at 37°C, recombinant individual clones were selected. Plasmid DNA was isolated from a transformant with the Qiaprep Spin Miniprep Kit (Product No. 27106, Qiagen, Hilden, Germany) in accordance with the manufacturer's instructions and checked by restriction cleavage. The resulting plasmid was called pCREmetAEY. It is shown in figure 2. The strain E. coli DH5 α mcrr/pCREmetAEY was deposited as a pure culture on 14th June 2001 at the Deutsche Sammlung für Mikroorganismen und Zellkulturen (DSMZ = German Collection of Microorganisms and Cell Cultures, Braunschweig, Germany) in accordance with the Budapest Treaty as DSM 14353.

5.4 Preparation of the strain C. glutamicum ATCC13032/pCREmetAEY

The vector pCREmetAY obtained in example 5.3 was electroporated in the strain C. glutamicum ATCC13032 using the electroporation method described by Liebl et al. (FEMS Microbiology Letters, 53:299-303 (1989)). Selection of

plasmid-carrying cells took place on LBHIS agar comprising 18.5 g/l brain-heart infusion broth, 0.5 M sorbitol, 5 g/l Bacto-tryptone, 2.5 g/l Bacto-yeast extract, 5 g/l NaCl and 18 g/l Bacto-agar, which had been supplemented with 25 mg/l kanamycin. Incubation was carried out for 2 days at 33°C.

Plasmid DNA was isolated from a transformant by conventional methods (Peters-Wendisch et al., 1998, Microbiology 144, 915-927) and checked by restriction cleavage. The resulting strain was called ATCC13032pCREmetAEY.

10 Example 6

Fermentative preparation of L-methionine with the strain ATCC13032/pCREmetAEY

The strain *C. glutamicum* ATCC13032/pCREmetAEY constructed by the process described in example 4 was cultured in a nutrient medium suitable for the production of methionine and the methionine content in the culture supernatant was determined.

For this, the strain was first incubated on an agar plate with the corresponding antibiotic (brain-heart agar with kanamycin (25 mg/l)) for 24 hours at 33°C. Starting from this agar plate culture, a preculture was seeded (10 ml medium in a 100 ml conical flask). The medium MM-1 was used as the medium for the preculture.

Medium MM-1

CSL (corn steep liquor) 5 g/l

MOPS (morpholinopropanesulfonic acid) 20 g/l

Glucose (autoclaved separately) 50g/l

Salts:

(NH₄)₂SO₄ 25 g/l

KH₂PO₄ 0.1 g/l

MgSO ₄ * 7 H ₂ O	1.0 g/l
CaCl ₂ * 2 H ₂ O	10 mg/l
FeSO ₄ * 7 H ₂ O	10 mg/l
MnSO ₄ * H ₂ O	5.0mg/l
Biotin (sterile-filtered)	0.01 mg/l
Vitamin B12 (sterile-filtered)	0.02 mg/l
Thiamine * HCl (sterile-filtered)	0.2 mg/l
CaCO ₃	25 g/l

The CSL, MOPS and the salt solution were brought to pH 7 with aqueous ammonia and autoclaved. The sterile substrate and vitamin solutions were then added, as well as the CaCO₃ autoclaved in the dry state. Kanamycin (25 mg/l) was added to this.

The preculture was incubated for 16 hours at 33°C at 240 rpm on a shaking machine and then used as the inoculum for the main culture in the fermenter. To establish an optical density(at 660 nm) of 1.0 as the starting value for the main culture in the fermenter, the corresponding amount of culture broth was transferred from the preculture.

The medium MM-2, which has the following composition, was used for the main culture:

Medium MM-2

CSL (corn steep liquor)	5 g/l
Glucose (autoclaved separately)	50g/l
Salts:	
(NH ₄) ₂ SO ₄	25 g/l
KH ₂ PO ₄	0.1 g/l

MgSO ₄ * 7 H ₂ O	1.0 g/l
CaCl ₂ * 2 H ₂ O	10 mg/l
FeSO ₄ * 7 H ₂ O	10 mg/l
MnSO ₄ * H ₂ O	5.0mg/l
Biotin (sterile-filtered)	0.01 mg/l
Vitamin B12 (sterile-filtered)	0.02 mg/l
Thiamine * HCl (sterile-filtered)	0.2 mg/l
Antifoam (Structol)	0.5 g/l

All the components of the medium were initially introduced directly into the fermenter, dissolved in water and then sterilized by means of heat (121°C, 20 minutes). Only the glucose was prepared in a stock solution of 50 wt.% and sterilized separately (also 121°C, 20 minutes). Biotin or thiamine were sterile-filtered and added under aseptic conditions directly before the start of fermentation.

Culturing was carried out by the batch process in a bioreactor with a working volume of 0.5 L (Multifermenter SIXFORS from Infors GmbH, Bodmingen, Switzerland). After addition of the inoculum, the starting volume in the fermenter was 0.4 L in total. Further culturing was carried out under constant aeration (0.1 vvm ("volume per volume per minute") and stirring at 33°C and a pH of 7.0. Correction or adjustment of the pH was carried out with a 5% NH₄OH solution. The set value for the concentration of dissolved oxygen in the fermentation medium was regulated at 40% and adjusted via the stirrer speed at a constant rate of aeration.

After 48 hours the process was ended and the optical density (OD) of the culture suspension was determined with an LP2W photometer from Dr. Lange (Berlin, Germany) at a measurement wavelength of 660 nm. The concentration of L-methionine formed was determined with an amino acid analyzer from Eppendorf-

BioTronik (Hamburg, Germany) by ion exchange chromatography and post-column derivation with ninhydrin detection.

5 An optical density in the final sample of 31.7 and a concentration of L-methionine of 39.0 mg per liter could be determined by the methods described above as the result.

Example 7

Preparation of biomass-free broth containing L-methionine

10 The biomass was first separated off from a fermentation broth comprising L-methionine prepared by the process of example 6 and comprising about 39 mg/l L-methionine. For this, 0.5 l of the above-mentioned fermentation broth was centrifuged with a laboratory centrifuge of the Biofuge-Stratos type from Heraeus (Düsseldorf, Germany) for 20 minutes at 4,000 rpm and the supernatant from the centrifugation was then purified further by cross-flow ultrafiltration with an MRC polymer membrane of 30kD in an ultrafiltrations unit from ICT GmbH (Bad Homburg, Germany).

Example 8

20 Preparation of a biomass-free product comprising L-methionine from a fermentation broth

25 The biomass was first separated off from a fermentation broth comprising L-methionine prepared by the process as described under example 6 and comprising about 39.0 mg/l L-methionine. For this, the fermenter contents of the above-mentioned fermentation broth were centrifuged and subjected to ultrafiltration as described in example 7.

30 23.7 g pure L-methionine (>99%; MERCK, Darmstadt, Germany) were then added batchwise to 300 g of the biomass-free filtrate, while stirring, in order to establish the desired content of L-methionine in the product. The suspension comprising L-methionine treated in this way was then mixed with 150 g water, with further stirring, to improve the working-up properties.

5 A portion of the suspension improved in this way was then lyophilized in a freeze-dryer of the type LYOVAC GT 2 from Leybold (Cologne, Germany). The product comprising L-methionine prepared in this manner had a content of 70 wt.% L-methionine and was free-flowing.

10 The remaining portion of the suspension improved in this way was treated by means of spray drying in a laboratory spray dryer of the Büchi-190 type from Büchi-Labortechnik GmbH (Constance, Germany) at an intake temperature of 170°C, a starting temperature of 105°C, a pressure difference of -40 mbar and an air flow rate of 600 NL/h. The product comprising L-methionine prepared in this manner had a content of 70 wt.% L-methionine and was free-flowing.

Example 9

15 Preparation of a biomass-containing product comprising L-methionine from a fermentation broth

20 From a fermentation broth comprising L-methionine prepared by the process of example 6 and comprising about 39 mg/l L-methionine, 23.7 g pure L-methionine (>99%; MERCK, Darmstadt, Germany) was first added batchwise, while stirring, in order to establish the desired content of L-methionine in the product. The fermentation broth treated in this way was then mixed with 150 g water, with further stirring, to improve the working-up properties.

25 A portion of this biomass-containing broth was then lyophilized in a freeze-dryer of the type LYOVAC GT 2 from Leybold (Cologne, Germany). The product comprising L-methionine prepared in this way had a content of 65 wt.% L-methionine and was free-flowing.

30 The remaining portion of the biomass-containing broth was treated by means of spray drying in a laboratory spray dryer of the Büchi-190 type from Büchi-Labortechnik GmbH (Constance, Germany) at an intake temperature of 170°C, a starting temperature of 105°C, a pressure difference of -40 mbar and an

air flow rate of 600 NL/h. The product comprising L-methionine prepared in this way had a content of 65 wt.% L-methionine and was free-flowing.

Brief Description of the Figures:

- Figure 1: Plasmid pCREmetAE
- Figure 2: pCREmetAEY

The abbreviations used in the figures have the following meaning:

Km: Resistance gene for kanamycin

metE: metE gene of *C. glutamicum*

metY: metY gene of *C. glutamicum*

metA: metA gene of *C. glutamicum*

Ptac: tac promoter

rrnB-T1T2: Terminator T1T2 of the rrnB gene of *E. coli*

rep: Plasmid-coded replication origin for *C. glutamicum*
(of pHM1519)

BamHI: Cleavage site of the restriction enzyme BamHI

EcoRI: Cleavage site of the restriction enzyme EcoRI

SalI: Cleavage site of the restriction enzyme SalI

This disclosure is based on priority documents DE 100 38 023.9, DE 101 09 689.5 and US 60/294,250, each incorporated by reference.

Obviously, numerous modifications of the invention are possible in view of the above teachings. Therefore, within the scope of the appended claims, the invention may be practiced otherwise than as specifically described herein.

SEQUENCE PROTOCOL

<110> Degussa AG

5 <120> Nucleotide sequences which code for the metE gene

<130> 000361 BT

<140>

10 <141>

<160> 8

<170> PatentIn Ver. 2.1

15

<210> 1

<211> 2810

<212> DNA

<213> Corynebacterium glutamicum

20

<220>

<221> CDS

<222> (317)..(2551)

<223> metE gene

25

<400> 1

agcccaaaac ggcaccatga atttaaattcc ccggaacttc ttgacagacc gagcagtcta 60

ggggtttgggtt gaaaacgcaa tcggttcact tttaatcctc tccctggagc cccggatgat 120

30

gaggaacgcc aaagctttct gaatggaaat tttaagcgtt aagtgggacg acctcgatta 180

caaaaaggcg aggaaacccc cggggcagct ttctgccacc cgggtgatttc gcgaaccttg 240

35

aaacatcgtc agaagattgc cgtgcgtcct agccgggatc cgcacgttcg gctcaagcag 300

aaagtcttta actcac atg act tcc aac ttt tct tcc act gtc gct ggt ctt 352

Met Thr Ser Asn Phe Ser Ser Thr Val Ala Gly Leu
1 5 10

40

cct cgc atc gga gcg aag cgt gaa ctg aag ttc gcg ctc gaa ggc tac 400

Pro Arg Ile Gly Ala Lys Arg Glu Leu Lys Phe Ala Leu Glu Gly Tyr
15 20 25

45

tgg aat gga tca att gaa ggt cgc gaa ctt gcg cag acc gcc cgc caa 448

Trp Asn Gly Ser Ile Glu Gly Arg Glu Leu Ala Gln Thr Ala Arg Gln
30 35 40

50

ttg gtc aac act gca tcg gat tct ttg tct gga ttg gat tcc gtt ccg 496

Leu Val Asn Thr Ala Ser Asp Ser Leu Ser Gly Leu Asp Ser Val Pro
45 50 55 60

55

ttt gca gga cgt tcc tac tac gac gca atg ctc gat acc gcc gct att 544

Phe Ala Gly Arg Ser Tyr Tyr Asp Ala Met Leu Asp Thr Ala Ala Ile
65 70 75

60

ttg ggt gtg ctg ccg gag cgt ttt gat gac atc gct gat cat gaa aac 592

Leu Gly Val Leu Pro Glu Arg Phe Asp Asp Ile Ala Asp His Glu Asn
80 85 90

gat ggt ctc cca ctg tgg att gac cgc tac ttt ggc gct gct cgc ggt 640

	Asp	Gly	Leu	Pro	Leu	Trp	Ile	Asp	Arg	Tyr	Phe	Gly	Ala	Ala	Arg	Gly	
			95					100					105				
5	act	gag	acc	ctg	cct	gca	cag	gca	atg	acc	aag	tgg	ttt	gat	acc	aac	688
	Thr	Glu	Thr	Leu	Pro	Ala	Gln	Ala	Met	Thr	Lys	Trp	Phe	Asp	Thr	Asn	
		110					115					120					
10	tac	cac	tac	ctc	gtg	ccg	gag	ttg	tct	gcg	gat	aca	cgt	ttc	gtt	ttg	736
	Tyr	His	Tyr	Leu	Val	Pro	Glu	Leu	Ser	Ala	Asp	Thr	Arg	Phe	Val	Leu	
		125				130					135					140	
15	gat	gcg	tcc	gcg	ctg	att	gag	gat	ctc	cgt	tgc	cag	cag	gtt	cgt	ggc	784
	Asp	Ala	Ser	Ala	Leu	Ile	Glu	Asp	Leu	Arg	Cys	Gln	Gln	Val	Arg	Gly	
					145					150					155		
20	gtt	aat	gcc	cgc	cct	gtt	ctg	gtt	ggc	cca	ctg	act	ttc	ctt	tcc	ctt	832
	Val	Asn	Ala	Arg	Pro	Val	Leu	Val	Gly	Pro	Leu	Thr	Phe	Leu	Ser	Leu	
				160					165					170			
25	gct	cgc	acc	act	gat	ggc	tcc	aat	cct	ttg	gat	cac	ctg	cct	gca	ctg	880
	Ala	Arg	Thr	Thr	Asp	Gly	Ser	Asn	Pro	Leu	Asp	His	Leu	Pro	Ala	Leu	
			175					180					185				
30	ttt	gag	gtc	tac	gag	cgc	ctc	atc	aag	tct	ttc	gat	act	gag	tgg	gtt	928
	Phe	Glu	Val	Tyr	Glu	Arg	Leu	Ile	Lys	Ser	Phe	Asp	Thr	Glu	Trp	Val	
		190					195					200					
35	cag	atc	gat	gag	cct	gcg	ttg	gtc	acc	gat	gtt	gct	cct	gag	gtt	ttg	976
	Gln	Ile	Asp	Glu	Pro	Ala	Leu	Val	Thr	Asp	Val	Ala	Pro	Glu	Val	Leu	
		205				210					215					220	
40	gag	cag	gtc	cgc	gct	ggc	tac	acc	act	ttg	gct	aag	cgc	gat	ggc	gtg	1024
	Glu	Gln	Val	Arg	Ala	Gly	Tyr	Thr	Thr	Leu	Ala	Lys	Arg	Asp	Gly	Val	
				225						230					235		
45	ttt	gtc	aat	act	tac	ttc	ggc	tct	ggc	gat	cag	gcg	ctg	aac	act	ctt	1072
	Phe	Val	Asn	Thr	Tyr	Phe	Gly	Ser	Gly	Asp	Gln	Ala	Leu	Asn	Thr	Leu	
				240				245						250			
50	gcg	ggc	atc	ggc	ctt	ggc	gcg	att	ggc	gtt	gac	ttg	gtc	acc	cat	ggc	1120
	Ala	Gly	Ile	Gly	Leu	Gly	Ala	Ile	Gly	Val	Asp	Leu	Val	Thr	His	Gly	
			255				260						265				
55	gtc	act	gag	ctt	gct	gcg	tgg	aag	ggc	gag	gag	ctg	ctg	gtt	gcg	ggc	1168
	Val	Thr	Glu	Leu	Ala	Ala	Trp	Lys	Gly	Glu	Glu	Leu	Leu	Val	Ala	Gly	
		270					275					280					
60	atc	gtt	gat	ggc	cgt	aac	att	tgg	cgc	acc	gac	ctg	tgt	gct	gct	ctt	1216
	Ile	Val	Asp	Gly	Arg	Asn	Ile	Trp	Arg	Thr	Asp	Leu	Cys	Ala	Ala	Leu	
		285				290					295					300	
65	gct	tcc	ctg	aag	cgc	ctg	gca	gct	cgc	ggc	cca	atc	gca	gtg	tct	acc	1264
	Ala	Ser	Leu	Lys	Arg	Leu	Ala	Ala	Arg	Gly	Pro	Ile	Ala	Val	Ser	Thr	
					305					310					315		

	tct	tgt	tca	ctg	ctg	cac	gtt	cct	tac	acc	ctc	gag	gct	gag	aac	att	1312
	Ser	Cys	Ser	Leu	Leu	His	Val	Pro	Tyr	Thr	Leu	Glu	Ala	Glu	Asn	Ile	
				320					325					330			
5	gag	cct	gag	gtc	cgc	gac	tgg	ctt	gcc	ttc	ggc	tgc	gag	aag	atc	acc	1360
	Glu	Pro	Glu	Val	Arg	Asp	Trp	Leu	Ala	Phe	Gly	Ser	Glu	Lys	Ile	Thr	
				335				340					345				
10	gag	gtc	aag	ctg	ctt	gcc	gac	gcc	cta	gcc	ggc	aac	atc	gac	gcg	gct	1408
	Glu	Val	Lys	Leu	Leu	Ala	Asp	Ala	Leu	Ala	Gly	Asn	Ile	Asp	Ala	Ala	
				350				355					360				
15	gcg	ttc	gat	gcg	gcg	tcc	gca	gca	att	gct	tct	cga	cgc	acc	tcc	cca	1456
	Ala	Phe	Asp	Ala	Ala	Ser	Ala	Ala	Ile	Ala	Ser	Arg	Arg	Thr	Ser	Pro	
							370					375				380	
	cgc	acc	gca	cca	atc	acg	cag	gaa	ctc	cct	ggc	cgt	agc	cgt	gga	tcc	1504
	Arg	Thr	Ala	Pro	Ile	Thr	Gln	Glu	Leu	Pro	Gly	Arg	Ser	Arg	Gly	Ser	
							385				390				395		
20	ttc	gac	act	cgt	gtt	acg	ctg	cag	gag	aag	tca	ctg	gag	ctt	cca	gct	1552
	Phe	Asp	Thr	Arg	Val	Thr	Leu	Gln	Glu	Lys	Ser	Leu	Glu	Leu	Pro	Ala	
							400							410			
25	ctg	cca	acc	acc	acc	att	ggc	tct	ttc	cca	cag	acc	cca	tcc	att	cgt	1600
	Leu	Pro	Thr	Thr	Thr	Ile	Gly	Ser	Phe	Pro	Gln	Thr	Pro	Ser	Ile	Arg	
								420					425				
30	tct	gct	cgc	gct	cgt	ctg	cgc	aag	gaa	tcc	atc	act	ttg	gag	cag	tac	1648
	Ser	Ala	Arg	Ala	Arg	Leu	Arg	Lys	Glu	Ser	Ile	Thr	Leu	Glu	Gln	Tyr	
								435					440				
35	gaa	gag	gca	atg	cgc	gaa	gaa	atc	gat	ctg	gtc	atc	gcc	aag	cag	gaa	1696
	Glu	Glu	Ala	Met	Arg	Glu	Glu	Ile	Asp	Leu	Val	Ile	Ala	Lys	Gln	Glu	
								450			455					460	
40	gaa	ctt	ggc	ctt	gat	gtg	ttg	gtt	cac	ggc	gag	cca	gag	cgc	aac	gac	1744
	Glu	Leu	Gly	Leu	Asp	Val	Leu	Val	His	Gly	Glu	Pro	Glu	Arg	Asn	Asp	
										470					475		
45	atg	gtt	cag	tac	ttc	tct	gaa	ctt	ctc	gac	ggc	ttc	ctc	tca	acc	gcc	1792
	Met	Val	Gln	Tyr	Phe	Ser	Glu	Leu	Leu	Asp	Gly	Phe	Leu	Ser	Thr	Ala	
														490			
50	aac	ggc	tgg	gtc	caa	agc	tac	ggc	tcc	cgc	tgt	gtt	cgt	cct	cca	gtg	1840
	Asn	Gly	Trp	Val	Gln	Ser	Tyr	Gly	Ser	Arg	Cys	Val	Arg	Pro	Pro	Val	
								500					505				
55	ttg	ttc	gga	aac	gtt	tcc	cgc	cca	gcg	cca	atg	act	gtc	aag	tgg	ttc	1888
	Leu	Phe	Gly	Asn	Val	Ser	Arg	Pro	Ala	Pro	Met	Thr	Val	Lys	Trp	Phe	
								515					520				
60	cag	tac	gca	cag	agc	ctg	acc	cag	aag	cat	gtc	aag	gga	atg	ctc	acc	1936
	Gln	Tyr	Ala	Gln	Ser	Leu	Thr	Gln	Lys	His	Val	Lys	Gly	Met	Leu	Thr	
								530			535					540	
65	ggc	cca	gtc	acc	atc	ctt	gca	tgg	tcc	ttc	gtt	cgc	gat	gat	cag	ccg	1984
	Gly	Pro	Val	Thr	Ile	Leu	Ala	Trp	Ser	Phe	Val	Arg	Asp	Asp	Gln	Pro	
															555		
70	ctg	gct	acc	act	gct	gac	cag	gtt	gca	ctg	gca	ctg	cgc	gat	gaa	att	2032

Leu Ala Thr Thr Ala Asp Gln Val Ala Leu Ala Leu Arg Asp Glu Ile
 560 565 570
 5 aac gat ctc atc gag gct ggc gcg aag atc atc cag gtg gat gag cct 2080
 Asn Asp Leu Ile Glu Ala Gly Ala Lys Ile Ile Gln Val Asp Glu Pro
 575 580 585
 10 gcg att cgt gaa ctg ttg ccg cta cga gac gtc gat aag cct gcc tac 2128
 Ala Ile Arg Glu Leu Leu Pro Leu Arg Asp Val Asp Lys Pro Ala Tyr
 590 595 600
 15 ctg cag tgg tcc gtg gac tcc ttc cgc ctg gcg act gcc ggc gca ccc 2176
 Leu Gln Trp Ser Val Asp Ser Phe Arg Leu Ala Thr Ala Gly Ala Pro
 605 610 615 620
 gac gac gtc caa atc cac acc cac atg tgc tac tcc gag ttc aac gaa 2224
 Asp Asp Val Gln Ile His Thr His Met Cys Tyr Ser Glu Phe Asn Glu
 625 630 635
 20 gtg atc tcc tcg gtc atc gcg ttg gat gcc gat gtc acc acc atc gaa 2272
 Val Ile Ser Ser Val Ile Ala Leu Asp Ala Asp Val Thr Thr Ile Glu
 640 645 650
 gca gca cgt tcc gac atg cag gtc ctc gct gct ctg aaa tct tcc ggc 2320
 Ala Ala Arg Ser Asp Met Gln Val Leu Ala Ala Leu Lys Ser Ser Gly
 655 660 665
 25 ttc gag ctc ggc gtc gga cct ggt gtg tgg gat atc cac tcc ccg cgc 2368
 Phe Glu Leu Gly Val Gly Pro Gly Val Trp Asp Ile His Ser Pro Arg
 670 675 680
 gtt cct tcc gcg cag gaa gtg gac ggt ctc ctc gag gct gca ctg cag 2416
 Val Pro Ser Ala Gln Glu Val Asp Gly Leu Leu Glu Ala Ala Leu Gln
 685 690 695 700
 tcc gtg gat cct cgc cag ctg tgg gtc aac cca gac tgt ggt ctg aag 2464
 Ser Val Asp Pro Arg Gln Leu Trp Val Asn Pro Asp Cys Gly Leu Lys
 705 710 715
 40 acc cgt gga tgg cca gaa gtg gaa gct tcc cta aag gtt ctc gtt gag 2512
 Thr Arg Gly Trp Pro Glu Val Glu Ala Ser Leu Lys Val Leu Val Glu
 720 725 730
 45 tcc gct aag cag gct cgt gag aaa atc gga gca act atc taaattgggt 2561
 Ser Ala Lys Gln Ala Arg Glu Lys Ile Gly Ala Thr Ile
 735 740 745
 taccgctagg aacccaaaga ttaagggcac gagtgtcacc aggattgccg cacccatggc 2621
 50 aacaccgaag gacaccgtgc ccactcctat ttgcatcaca gcgccaagg tagcggcgcc 2681
 caaaacagcg cccacctggc gtgaggtgtt gtaaaaacca gaagcagagc ccactaaatc 2741
 ctgcggaaca tcacgcagag caatcacaga gttcggtgca aaactcatcg cgttggagct 2801
 55 accgaacaa 2810
 60 <210> 2
 <211> 745
 <212> PRT

<213> Corynebacterium glutamicum

<400> 2

5	Met	Thr	Ser	Asn	Phe	Ser	Ser	Thr	Val	Ala	Gly	Leu	Pro	Arg	Ile	Gly	1	5	10	15
	Ala	Lys	Arg	Glu	Leu	Lys	Phe	Ala	Leu	Glu	Gly	Tyr	Trp	Asn	Gly	Ser	20	25	30	
10	Ile	Glu	Gly	Arg	Glu	Leu	Ala	Gln	Thr	Ala	Arg	Gln	Leu	Val	Asn	Thr	35	40	45	
	Ala	Ser	Asp	Ser	Leu	Ser	Gly	Leu	Asp	Ser	Val	Pro	Phe	Ala	Gly	Arg	50	55	60	
15	Ser	Tyr	Tyr	Asp	Ala	Met	Leu	Asp	Thr	Ala	Ala	Ile	Leu	Gly	Val	Leu	65	70	75	80
	Pro	Glu	Arg	Phe	Asp	Asp	Ile	Ala	Asp	His	Glu	Asn	Asp	Gly	Leu	Pro	85	90	95	
20	Leu	Trp	Ile	Asp	Arg	Tyr	Phe	Gly	Ala	Ala	Arg	Gly	Thr	Glu	Thr	Leu	100	105	110	
25	Pro	Ala	Gln	Ala	Met	Thr	Lys	Trp	Phe	Asp	Thr	Asn	Tyr	His	Tyr	Leu	115	120	125	
	Val	Pro	Glu	Leu	Ser	Ala	Asp	Thr	Arg	Phe	Val	Leu	Asp	Ala	Ser	Ala	130	135	140	
	Leu	Ile	Glu	Asp	Leu	Arg	Cys	Gln	Gln	Val	Arg	Gly	Val	Asn	Ala	Arg	145	150	155	160
	Pro	Val	Leu	Val	Gly	Pro	Leu	Thr	Phe	Leu	Ser	Leu	Ala	Arg	Thr	Thr	165	170	175	
	Asp	Gly	Ser	Asn	Pro	Leu	Asp	His	Leu	Pro	Ala	Leu	Phe	Glu	Val	Tyr	180	185	190	
40	Glu	Arg	Leu	Ile	Lys	Ser	Phe	Asp	Thr	Glu	Trp	Val	Gln	Ile	Asp	Glu	195	200	205	
	Pro	Ala	Leu	Val	Thr	Asp	Val	Ala	Pro	Glu	Val	Leu	Glu	Gln	Val	Arg	210	215	220	
45	Ala	Gly	Tyr	Thr	Thr	Leu	Ala	Lys	Arg	Asp	Gly	Val	Phe	Val	Asn	Thr	225	230	235	240
	Tyr	Phe	Gly	Ser	Gly	Asp	Gln	Ala	Leu	Asn	Thr	Leu	Ala	Gly	Ile	Gly	245	250	255	
50	Leu	Gly	Ala	Ile	Gly	Val	Asp	Leu	Val	Thr	His	Gly	Val	Thr	Glu	Leu	260	265	270	

Ala Ala Trp Lys Gly Glu Glu Leu Val Ala Gly Ile Val Asp Gly
 275 280 285
 5 Arg Asn Ile Trp Arg Thr Asp Leu Cys Ala Ala Leu Ala Ser Leu Lys
 290 295 300
 Arg Leu Ala Ala Arg Gly Pro Ile Ala Val Ser Thr Ser Cys Ser Leu
 305 310 315 320
 10 Leu His Val Pro Tyr Thr Leu Glu Ala Glu Asn Ile Glu Pro Glu Val
 325 330 335
 Arg Asp Trp Leu Ala Phe Gly Ser Glu Lys Ile Thr Glu Val Lys Leu
 340 345 350
 15 Leu Ala Asp Ala Leu Ala Gly Asn Ile Asp Ala Ala Ala Phe Asp Ala
 355 360 365
 Ala Ser Ala Ala Ile Ala Ser Arg Arg Thr Ser Pro Arg Thr Ala Pro
 370 375 380
 20 Ile Thr Gln Glu Leu Pro Gly Arg Ser Arg Gly Ser Phe Asp Thr Arg
 385 390 395 400
 25 Val Thr Leu Gln Glu Lys Ser Leu Glu Leu Pro Ala Leu Pro Thr Thr
 405 410 415
 Thr Ile Gly Ser Phe Pro Gln Thr Pro Ser Ile Arg Ser Ala Arg Ala
 420 425 430
 30 Arg Leu Arg Lys Glu Ser Ile Thr Leu Glu Gln Tyr Glu Glu Ala Met
 435 440 445
 Arg Glu Glu Ile Asp Leu Val Ile Ala Lys Gln Glu Glu Leu Gly Leu
 450 455 460
 35 Asp Val Leu Val His Gly Glu Pro Glu Arg Asn Asp Met Val Gln Tyr
 465 470 475 480
 40 Phe Ser Glu Leu Leu Asp Gly Phe Leu Ser Thr Ala Asn Gly Trp Val
 485 490 495
 Gln Ser Tyr Gly Ser Arg Cys Val Arg Pro Pro Val Leu Phe Gly Asn
 500 505 510
 45 Val Ser Arg Pro Ala Pro Met Thr Val Lys Trp Phe Gln Tyr Ala Gln
 515 520 525
 Ser Leu Thr Gln Lys His Val Lys Gly Met Leu Thr Gly Pro Val Thr
 530 535 540
 Ile Leu Ala Trp Ser Phe Val Arg Asp Asp Gln Pro Leu Ala Thr Thr
 545 550 555 560
 55 Ala Asp Gln Val Ala Leu Ala Leu Arg Asp Glu Ile Asn Asp Leu Ile
 565 570 575
 Glu Ala Gly Ala Lys Ile Ile Gln Val Asp Glu Pro Ala Ile Arg Glu
 580 585 590
 60 Leu Leu Pro Leu Arg Asp Val Asp Lys Pro Ala Tyr Leu Gln Trp Ser

595 600 605

Val Asp Ser Phe Arg Leu Ala Thr Ala Gly Ala Pro Asp Asp Val Gln
610 615 620

5 Ile His Thr His Met Cys Tyr Ser Glu Phe Asn Glu Val Ile Ser Ser
625 630 635 640

10 Val Ile Ala Leu Asp Ala Asp Val Thr Thr Ile Glu Ala Ala Arg Ser
645 650 655

Asp Met Gln Val Leu Ala Ala Leu Lys Ser Ser Gly Phe Glu Leu Gly
660 665 670

15 Val Gly Pro Gly Val Trp Asp Ile His Ser Pro Arg Val Pro Ser Ala
675 680 685

Gln Glu Val Asp Gly Leu Leu Glu Ala Ala Leu Gln Ser Val Asp Pro
690 695 700

20 Arg Gln Leu Trp Val Asn Pro Asp Cys Gly Leu Lys Thr Arg Gly Trp
705 710 715 720

25 Pro Glu Val Glu Ala Ser Leu Lys Val Leu Val Glu Ser Ala Lys Gln
725 730 735

Ala Arg Glu Lys Ile Gly Ala Thr Ile
740 745

30

35 <210> 3
<211> 41
<212> DNA
<213> Artificial sequence

40 <220>
<223> Description of the artificial sequence: Primer
metA-EVP5

45 <400> 3
agaacgaatt caaaggagga caaccatgcc caccctcgcg c 41

50 <210> 4
<211> 31
<212> DNA
<213> Künstliche Sequenz

55 <220>
<223> Beschreibung der künstlichen Sequenz: Primer
metA-EVP3

60 <400> 4
gtcgtggatc ccctattaga tgtagaactc g 31

<210> 5
<211> 47
<212> DNA
<213> Artificial sequence

5 <220>
 <223> Description of the artificial sequence: Primer
 metE-EVP5
 <400> 5
 ggctcaaaga tctaaaggag gacaaccatg acttccaact tttcttc 47

10 <210> 6
 <211> 40
 <212> DNA
 <213> Artificial sequence

15 <220>
 <223> Description of the artificial sequence: Primer
 metE-EVP3
 <400> 6
 20 ggttcctgtc gacggtacca tttagatagt tgctccgatt 40

25 <210> 7
 <211> 42
 <212> DNA
 <213> Artificial sequence

30 <220>
 <223> Description of the artificial sequence: Primer
 metY-EVP5
 <400> 7
 35 ctaataagtc gacaaaggag gacaaccatg ccaaagtacg ac 42

40 <210> 8
 <211> 33
 <212> DNA
 <213> Artificial sequence

45 <220>
 <223> Description of the artificial sequence: Primer
 metE-EVP3
 <400> 8
 gagtctaag catgctagat tgcagcaaag ccg 33